

4D08: MECHANISM OF IGNITION BY NON-EQUILIBRIUM PLASMA.

*Nikolay L. Aleksandrov, Svetlana V. Kindysheva, Ilya N. Kosarev,
Svetlana M. Starikovskaia, Andrey Yu. Starikovskii
MIPT, Russia*

Comment Godfrey Mungal, Stanford University, USA

mungal@stanford.edu

How do your conclusions for methane compare with those of [1], namely the pulsed plasma produces radicals which quickly decay leaving behind H₂ and CO which enhance flame ignition.

Reference:

[1] W. Kim, M.G. Mungal, M.A. Cappelli, *Appl. Phys. Letters*, 92, 051503 (2008) 3.

Reply by Andrey Starikovskiy

andreistarikovskii@gmail.com

We have demonstrated in [1,2] that for temperatures above the self-ignition threshold and high E/n values the discharge phase leads to atomic oxygen, atomic hydrogen and CH_x radicals formation. These radicals start the chemical chains and partially convert to CO and H₂. But CO and H₂ formation does not stop the chain ignition mechanism. Below self-ignition threshold CO and H₂ production competes with formation of peroxides, H₂O₂ and HO₂. These processes lead to mixture sensitization which enhances the ignition in longer time scale. For low temperature conditions we have strong competition between recombination and hydroperoxides formation due to reactions of hydrocarbons with O(1D) and OH radicals [3]. Thus we have different mechanisms of plasma assisted ignition for different initial temperatures.

References:

[1] Evgeny I. Mintoussov, Svetlana E. Yankina, Andrey A. Nikipelov, Svetlana S. Starikovskaia, Andrey Yu. Starikovskii, 45th AIAA Aerospace Sciences Meeting and Exhibit, 8–11 Jan 2007 Grand Sierra Resort Hotel Reno, Nevada. Paper AIAA-2007-1354.

[2] I.N. Kosarev, S.V. Kindusheva, N.L. Aleksandrov, S.M. Starikovskaia, A.Yu. Starikovskii, 45th AIAA Aerospace Sciences Meeting and Exhibit, 8–11 Jan 2007 Grand Sierra Resort Hotel Reno, Nevada. Paper AIAA-2007-1386.

[3] N.B. Anikin, S.M. Starikovskaia, A.Yu. Starikovskii, *Journal of Physics D: Applied Physics* 39 (2006) 3244–3252.